

# MÖBIUS FUNCTION OF COORDINATE HYPERPLANES IN COMPLEX ELLIPSOIDS

WITOLD JARNICKI

ABSTRACT. For  $p_1, \dots, p_n > 0$ , let  $\mathbb{E} = \{z \in \mathbb{C}^n : \sum_{j=1}^n |z_j|^{2p_j} < 1\}$  be a complex ellipsoid. We present effective formulas for the generalized Möbius and Green functions  $m_{\mathbb{E}}(A, \cdot)$ ,  $g_{\mathbb{E}}(A, \cdot)$  in the case where  $A := \{z \in \mathbb{E} : z_1 \cdots z_k = 0\}$  ( $1 \leq k \leq n$ ).

## 1. INTRODUCTION AND MAIN RESULTS

Let  $G \subset \mathbb{C}^n$  be a domain and let  $A \subset G$ . Define the *generalized Green function*

$$g_G(A, z) := \sup\{u(z) : u : G \longrightarrow [0, 1), \log u \in \mathcal{PSH}(G), \\ \forall_{a \in A} \exists_{C=C(u,a)>0} \forall_{w \in G} : u(w) \leq C\|w - a\|\}, \quad z \in G,$$

and the *generalized Möbius function*

$$m_G(A, z) := \sup\{|f(z)| : f \in \mathcal{O}(G, \mathbb{D}), f|_A \equiv 0\}, \quad z \in G,$$

where  $\mathbb{D}$  denotes the unit disc (cf. [Jar-Jar-Pfl 2003]).

The generalized Möbius and Green functions were recently studied by many authors (e.g. [Lár-Sig 1998]). Despite various properties proven, effective formulas for  $m_G(A, \cdot)$  and  $g_G(A, \cdot)$  are known only in a few special cases. An effective formula for the generalized Green function in the case of Euclidean ball  $G = \mathbb{B}_2 \subset \mathbb{C}^2$  and union of coordinate lines  $A = \{(z, w) \in \mathbb{B}_2 : zw = 0\}$  was given in [Ngu 2003].

Let  $n \geq 2$ . Let  $p_1, \dots, p_n > 0$ . Define the *complex ellipsoid with exponents*  $p_1, \dots, p_n$ :

$$\mathbb{E} = \mathbb{E}_{p_1, \dots, p_n} := \left\{z \in \mathbb{C}^n : \sum_{j=1}^n |z_j|^{2p_j} < 1\right\}.$$

Observe that  $\mathbb{E}_{p_1, \dots, p_n}$  is convex if and only if  $p_j \geq 1/2$ ,  $j = 1, \dots, n$  (cf. [Jar-Pfl 1993], § 8.4).

For  $G \subset \mathbb{C}^n$ ,  $k \in \{1, \dots, n\}$ , we consider the sets

$$A = A_{G,k} := \{z \in G : z_1 \cdots z_k = 0\}.$$

We give an effective formula for  $g_{\mathbb{E}}(A_{\mathbb{E},k}, \cdot)$  for any complex ellipsoid  $\mathbb{E} \subset \mathbb{C}^n$  and any  $k \in \{1, \dots, n\}$  (Theorem 1.2(a)). This is a generalization of a result in [Ngu 2003] (with a much simpler proof).

---

1991 *Mathematics Subject Classification.* 32F45, 32U35.

The author was supported in part by KBN grant no. 2 P03A 015 22.

We prove that the  $m_{\mathbb{E}}(A_{\mathbb{E},k}, \cdot) \equiv g_{\mathbb{E}}(A_{\mathbb{E},k}, \cdot)$  for the cases where  $\mathbb{E}$  is convex (Theorem 1.2(b)) or where  $k = 1$ ,  $n = 2$ ,  $p_2 \geq 1/2$  (Theorem 1.2(c)). This is not true in general nonconvex case; we prove that  $m_{\mathbb{E}}(A_{\mathbb{E},k}, \cdot) \not\equiv g_{\mathbb{E}}(A_{\mathbb{E},k}, \cdot)$  if there exists a  $j \in \{k+1, \dots, n\}$  such that  $p_j < 1/2$  (Theorem 1.2(d)). Partial results for the case  $k = n = 2$  are given (Theorem 1.2(e)).

**Definition 1.1.** Let  $p_1, \dots, p_n > 0$ ,  $1 \leq k \leq n$ . Put  $\mathbb{E} := \mathbb{E}_{p_1, \dots, p_n}$ ,  $A := A_{\mathbb{E},k}$ . Let  $z \in \mathbb{E}$  be such that the sequence  $(p_j |z_j|^{2p_j})_{j=1}^k$  is monotonically increasing. For  $s \in \{1, \dots, k\}$  define

$$q_s := \sum_{j=1}^s (2p_j)^{-1}, \quad r_s(z) := 1 - \sum_{j=s+1}^n |z_j|^{2p_j}, \quad c_s(z) := r_s(z)/q_s.$$

Let  $d := \max\{s \in \{1, \dots, k\} : 2p_s |z_s|^{2p_s} \leq c_s(z)\}$ . Define

$$(*) \quad R_{\mathbb{E}}(A, z) = \prod_{j=1}^d |z_j| \left( \frac{2p_j}{c_d(z)} \right)^{\frac{1}{2p_j}}.$$

The main results of the paper are the following:

**Theorem 1.2.** *Under the above assumptions we have:*

- (a)  $g_{\mathbb{E}}(A, \cdot) \equiv R_{\mathbb{E}}(A, \cdot)$ ,
- (b)  $m_{\mathbb{E}}(A, \cdot) \equiv g_{\mathbb{E}}(A, \cdot) \equiv R_{\mathbb{E}}(A, \cdot)$ , for  $p_j \geq 1/2$ ,  $j = 1, \dots, n$ ,
- (c)  $m_{\mathbb{E}}(A, \cdot) \equiv g_{\mathbb{E}}(A, \cdot) \equiv R_{\mathbb{E}}(A, \cdot)$ , for  $k = 1$ ,  $n = 2$ ,  $p_2 \geq 1/2$ ,
- (d)  $m_{\mathbb{E}}(A, \cdot) \not\equiv g_{\mathbb{E}}(A, \cdot)$ , if there exists a  $j \in \{k+1, \dots, n\}$  with  $p_j < 1/2$ .
- (e)  $m_{\mathbb{E}}(A, \cdot) \equiv g_{\mathbb{E}}(A, \cdot) \equiv R_{\mathbb{E}}(A, \cdot)$ , for  $k = n = 2$ ,  $p_1 \leq p_2$ , and either  $p_2 \geq 1/2$  or  $8p_1 + 4p_2(1 - p_2) > 1$ .

Observe that one can obtain the monotonicity of  $(p_j |z_j|^{2p_j})_{j=1}^k$  for arbitrary  $z \in \mathbb{E}$  by permutation of coordinates. The final result is a subdivision of  $\mathbb{E}$  into  $2^k - 1$  subsets, each with  $R_{\mathbb{E}}(A, \cdot)$  given by a formula of type (\*).

**Remark 1.3.** In the case where  $p_1 = \dots = p_n = 1$ , the domain  $\mathbb{E}$  is the Euclidean ball  $\mathbb{B}_n$ . Theorem 1.2 may then be formulated in the following simpler way.

**Corollary 1.4.** *Assume that  $|z_1| \leq \dots \leq |z_k|$  and let*

$$d := \max \left\{ s \in \{1, \dots, k\} : s|z_s|^2 + \sum_{j=s+1}^n |z_j|^2 \leq 1 \right\}.$$

*Then*

$$m_{\mathbb{B}_n}(A, z) = g_{\mathbb{B}_n}(A, z) = \left( \frac{d}{1 - \sum_{j=d+1}^n |z_j|^2} \right)^{\frac{d}{2}} \prod_{j=1}^d |z_j|.$$

**Remark 1.5.** It is unknown to the author whether  $m_{\mathbb{E}}(A, \cdot) \equiv g_{\mathbb{E}}(A, \cdot)$  for any  $p_j > 0$ ,  $j = 1, \dots, k$ ,  $p_j \geq 1/2$ ,  $j = k+1, \dots, n$ . However, Theorem 1.2(e) and author's research (cf. Remark 4.4) indicate that this is true at least in the case  $k = n = 2$ .

**Remark 1.6.** Take  $p_{j,\ell} \nearrow +\infty$ ,  $j = 1, \dots, n$ . Then

$$G_\ell := \mathbb{E}_{p_{1,\ell}, \dots, p_{n,\ell}} \nearrow \mathbb{D}^n.$$

Consequently,

$$m_{G_\ell}(A_{G_\ell, k}, \cdot) = g_{G_\ell}(A_{G_\ell, k}, \cdot) \searrow m_{\mathbb{D}^n}(A_{\mathbb{D}^n, k}, \cdot) = g_{\mathbb{D}^n}(A_{\mathbb{D}^n, k}, \cdot)$$

(cf. [Jar-Jar-Pfl 2003], Property 2.7) and hence

$$m_{\mathbb{D}^n}(A_{\mathbb{D}^n, k}, z) = |z_1| \cdots |z_k|, \quad z \in \mathbb{D}^n.$$

However, we need the formula for  $g_{\mathbb{D}^n}(A_{\mathbb{D}^n, k}, \cdot)$  before we prove Theorem 1.2 (cf. Lemma 2.1).

## 2. PROOF OF THEOREM 1.2

**Lemma 2.1.** *Let  $n \in \mathbb{N}$ ,  $1 \leq k \leq n$ . Then*

$$m_{\mathbb{D}^n}(A_{\mathbb{D}^n, k}, z) = g_{\mathbb{D}^n}(A_{\mathbb{D}^n, k}, z) = |z_1| \cdots |z_k|, \quad z \in \mathbb{D}^n.$$

*Proof of Lemma 2.1.* (Due to P.Pflug.) obviously,

$$|z_1| \cdots |z_k| \leq m_{\mathbb{D}^n}(A_{\mathbb{D}^n, k}, z) \leq g_{\mathbb{D}^n}(A_{\mathbb{D}^n, k}, z).$$

It remains to prove that  $g_{\mathbb{D}^n}(A_{\mathbb{D}^n, k}, z) \leq |z_1| \cdots |z_k|$ .

It is enough to prove that for any  $1 \leq k \leq n$  and for any function  $u : \mathbb{D}^n \rightarrow [0, 1]$  such that  $\log u \in \mathcal{PSH}(\mathbb{D}^n)$  and  $u(z) \leq C(a)\|z - a\|$ ,  $a \in A_{\mathbb{D}^n, k}$ ,  $z \in \mathbb{D}^n$ , we have  $u(z) \leq |z_1| \cdots |z_k|$ . We proceed by induction on  $k$ .

For  $k = 1$  the inequality follows from the Schwarz type lemma for logarithmically subharmonic functions  $u(\cdot, z_2, \dots, z_n)$ ,  $z_2, \dots, z_n \in \mathbb{D}$ .

For  $k > 1$  we first apply the case  $k = 1$  and get  $u(z_1, \dots, z_n) \leq |z_1|$ ,  $z \in \mathbb{D}^n$ . Applying the inductive assumption to  $u(z_1, \cdot)/|z_1|$ ,  $z_1 \in \mathbb{D}$ , finishes the proof.  $\square$

Take a  $u : \mathbb{E} \rightarrow [0, 1]$  with  $\log u \in \mathcal{PSH}(\mathbb{E})$  and  $u(\zeta) \leq C(a)\|\zeta - a\|$ ,  $a \in A$ ,  $\zeta \in \mathbb{E}$ . Consider the mapping

$$\begin{aligned} \iota_z : \mathbb{D}^d \ni (\zeta_1, \dots, \zeta_d) \\ \longmapsto \left( \zeta_1 \left( \frac{c_d(z)}{2p_1} \right)^{\frac{1}{2p_1}}, \dots, \zeta_d \left( \frac{c_d(z)}{2p_d} \right)^{\frac{1}{2p_d}}, z_{d+1}, \dots, z_n \right) \in \mathbb{E}. \end{aligned}$$

Applying the holomorphic contractivity of the generalized Green function and Lemma 2.1 proves that  $m_{\mathbb{E}}(A, z) \leq g_{\mathbb{E}}(A, z) \leq R_{\mathbb{E}}(A, z)$ .

It remains to prove that  $m_{\mathbb{E}}(A, z) \geq R_{\mathbb{E}}(A, z)$  (resp.  $g_{\mathbb{E}}(A, z) \geq R_{\mathbb{E}}(A, z)$ ). In the case  $d = k = n$  it suffices to take

$$f(\zeta) := \prod_{j=1}^d \zeta_j \left( \frac{2p_j}{c_d(z)} \right)^{\frac{1}{2p_j}}.$$

Then  $f \in \mathcal{O}(\mathbb{E}, \mathbb{D})$  and  $|f(z)| = R_{\mathbb{E}}(A, z)$ .

It remains to prove Theorem 1.2 in the remaining cases. We may assume that  $z_j \neq 0$ ,  $j = 1, \dots, d$ . Consider the following lemma.

**Lemma 2.2.** *Define  $\mathbb{E}' := \mathbb{E}_{p_{d+1}, \dots, p_n}$ .*

- (a) *Let  $v : \mathbb{E}' \rightarrow [0, 1)$ ,  $v \not\equiv 0$  be such that  $\log v \in \mathcal{PSH}(\mathbb{E}')$  and  $v(\zeta) \leq |\zeta_j|$ ,  $\zeta \in \mathbb{E}'$ ,  $j = d+1, \dots, k$ , and the mapping*

$$\mathbb{E}' \ni (\zeta_{d+1}, \dots, \zeta_n) \mapsto v(\zeta_{d+1}, \dots, \zeta_n) r_d^{q_d}(\zeta)$$

*attains its maximum value  $M$  at  $(z_{d+1}, \dots, z_n)$ . Then  $g_{\mathbb{E}}(A, z) = R_{\mathbb{E}}(A, z)$ .*

- (b) *The following conditions are equivalent:*

- *There exists an  $h \in \mathcal{O}(\mathbb{E}')$ ,  $h \not\equiv 0$  such that  $h(\zeta) = 0$  for  $\zeta_{d+1} \cdots \zeta_k = 0$  and the mapping*

$$\mathbb{E}' \ni (\zeta_{d+1}, \dots, \zeta_n) \mapsto |h(\zeta_{d+1}, \dots, \zeta_n)| r_d^{q_d}(\zeta)$$

*attains its maximum value  $M$  at the point  $(z_{d+1}, \dots, z_n)$ ,*

- *$m_{\mathbb{E}}(A, z) = R_{\mathbb{E}}(A, z)$ .*

We present a proof in Section 3. The result above reduces the proof of Theorem 1.2 to the following propositions.

**Proposition 2.3.** *There exists a  $v$  as required in Lemma 2.2(a).*

**Proposition 2.4.** *Using notation of Theorem 1.2, assume that  $p_j \geq 1/2$ ,  $j = d+1, \dots, n$ . Then there exists an  $h \in \mathcal{O}(\mathbb{E}')$  as required in Lemma 2.2(b).*

**Proposition 2.5.** *Using notation of Theorem 1.2, assume that  $p_{k+1} < 1/2$ . Then one cannot find an  $h$  as required in Lemma 2.2(b) for  $|z_\ell| \neq 0$  small enough,  $\ell = 1, \dots, k+1$  and  $z_\ell = 0$ ,  $\ell = k+2, \dots, n$ .*

**Proposition 2.6.** *Using notation of Theorem 1.2, assume that  $n = k = 2$ . Additionally, assume that  $p_2 \geq 1/2$  or  $8p_1 + 4p_2(1 - p_2) > 1$ . Then there exists an  $h \in \mathcal{O}(\mathbb{D})$  as required in Lemma 2.2(b).*

### 3. PROOF OF LEMMA 2.2

(a) Put

$$u(\zeta_1, \dots, \zeta_n) := M^{-1} \left( \prod_{j=1}^d (2p_j)^{\frac{1}{2p_j}} |\zeta_j| \right) q_d^{q_d} v(\zeta_{d+1}, \dots, \zeta_n).$$

Obviously  $\log u \in \mathcal{PSH}(\mathbb{E})$  and  $u(\zeta) \leq C|\zeta_j| \leq C\|\zeta - a\|$ ,  $\zeta \in \mathbb{E}$ , whenever  $a_j = 0$  for some  $j \in \{1, \dots, k\}$ . For  $\zeta \in \mathbb{E}$  we have:

$$u(\zeta) \leq M^{-1} \left( \frac{\sum_{j=1}^d |\zeta_j|^{2p_j}}{q_d} \right)^{q_d} q_d^{q_d} v(\zeta_{d+1}, \dots, \zeta_n) < 1. \quad ^1$$

Consequently  $u : \mathbb{E} \longrightarrow [0, 1)$ . On the other hand:

$$u(z) = M^{-1} R_{\mathbb{E}}(A, z) r_d^{q_d}(z) v(z_{d+1}, \dots, z_n) = R_{\mathbb{E}}(A, z).$$

(b) Assume that such an  $h$  exists. Put

$$f(\zeta_1, \dots, \zeta_n) := M^{-1} \left( \prod_{j=1}^d (2p_j)^{\frac{1}{2p_j}} \zeta_j \right) q_d^{q_d} h(\zeta_{d+1}, \dots, \zeta_n).$$

Observe that  $f(\zeta) = 0$  for  $\zeta \in A$ . Similarly as in (a) we prove that  $|f(\zeta)| < 1$ ,  $\zeta \in \mathbb{E}$  and  $f(z) = R_{\mathbb{E}}(A, z)$ .

Assume now that  $m_{\mathbb{E}}(A, z) = R_{\mathbb{E}}(A, z)$ . Let  $f \in \mathcal{O}(\mathbb{E}, \mathbb{D})$  be such that  $f|_A \equiv 0$  and  $|f(z)| = R_{\mathbb{E}}(A, z)$  (cf. [Jar-Jar-Pfl 2003], Property 2.5). Put

$$h(\zeta) := \frac{\partial^d f}{\partial z_1 \dots \partial z_d}(0, \zeta).$$

By definition, we have  $h(\zeta) = 0$  for  $\zeta_{d+1} \dots \zeta_k = 0$ . Applying the Schwarz lemma to the mapping  $f \circ \iota_{\zeta}$ ,  $\zeta \in \mathbb{E}'$ , we get

$$|h(\zeta_{d+1}, \dots, \zeta_n)| r_d^{q_d}(\zeta) \leq \left( \prod_{j=1}^d (2p_j)^{\frac{1}{2p_j}} \right) q_d^{q_d}, \quad \zeta \in \mathbb{E}',$$

$$|h(z_{d+1}, \dots, z_n)| r_d^{q_d}(z) = \left( \prod_{j=1}^d (2p_j)^{\frac{1}{2p_j}} \right) q_d^{q_d}.$$

#### 4. PROOF OF PROPOSITIONS 2.3, 2.4, 2.5, AND 2.6

*Proof of Proposition 2.3.* We may assume that  $z_{d+1}, \dots, z_n \geq 0$ . Consider functions of the form

$$v_{\alpha}(\zeta_{d+1}, \dots, \zeta_n) = \left( \prod_{j=d+1}^k |\zeta_j|^{1+\alpha_j} \right) \left( \prod_{j=k+1}^n |\zeta_j|^{\alpha_j} \right),$$

where  $\alpha = (\alpha_{d+1}, \dots, \alpha_n)$ ,  $\alpha_{d+1}, \dots, \alpha_n \geq 0$ . Obviously  $v : \mathbb{E}' \longrightarrow [0, 1)$ ,  $\log v \in \mathcal{PSH}(\mathbb{E}')$ , and  $v(\zeta) \leq |\zeta_j| \leq \|\zeta - a\|$ ,  $\zeta \in \mathbb{E}'$ , whenever  $a_j = 0$  for some  $j \in \{d+1, \dots, k\}$ . Since  $v_{\alpha}(\zeta_{d+1}, \dots, \zeta_n) =$

---

<sup>1</sup>Let  $a_1, \dots, a_d \geq 0$ ,  $w_1, \dots, w_d > 0$ . Then

$$\prod_{j=1}^d a_j^{w_j} \leq \left( \frac{\sum_{j=1}^d w_j a_j}{\sum_{j=1}^d w_j} \right)^{\sum_{j=1}^d w_j}.$$

$v_\alpha(|\zeta_{d+1}|, \dots, |\zeta_n|)$ , it is enough to find an  $\alpha$  such that the function

$$\mathbb{E}' \cap \mathbb{R}_+^{n-d} \ni (t_{d+1}, \dots, t_n) \mapsto v_\alpha(t_{d+1}, \dots, t_n) r_d^{q_d}(t)$$

attains its maximum at  $(z_{d+1}, \dots, z_n)$ . Considering the partial (logarithmic) derivatives results in the following equations:

$$\begin{aligned} 0 &= 1 + \alpha_j - 2p_j q_d \frac{t_j^{2p_j}}{r_d(t)}, \quad j = d+1, \dots, k, \\ 0 &= \alpha_j - 2p_j q_d \frac{t_j^{2p_j}}{r_d(t)}, \quad j = k+1, \dots, n. \end{aligned}$$

These give formulas for  $\alpha_{d+1}, \dots, \alpha_n$  such that  $(z_{d+1}, \dots, z_n)$  is the common zero of the derivatives. To prove that there are no other points like this, consider a reformulation of the above equations:

$$\begin{aligned} r_d(t) &= \frac{2p_j q_d t_j^{2p_j}}{1 + \alpha_j}, \quad j = d+1, \dots, k, \\ r_d(t) &= \frac{2p_j q_d t_j^{2p_j}}{\alpha_j}, \quad j = k+1, \dots, n. \end{aligned}$$

The left side is decreasing in any of the variables, while the right sides are increasing. Thus, at most one common zero is allowed.

It remains to check, whether  $\alpha_j \geq 0$ ,  $j = d+1, \dots, n$ . Obviously, this is true for  $j = k+1, \dots, n$  and in the remaining cases we have:

$$\alpha_j = \frac{2p_j q_d t_j^{2p_j} - r_d(t)}{t_j r_d(t)} \geq 0,$$

since this is the way we have chosen  $d$ . □

*Proof of Proposition 2.4.* We may assume that  $z_{d+1}, \dots, z_n \geq 0$ . Consider functions of the form

$$h_\alpha(\zeta_{d+1}, \dots, \zeta_n) = \left( \prod_{j=d+1}^k \zeta_j e^{\alpha_j \zeta_j} \right) \left( \prod_{j=k+1}^n e^{\alpha_j \zeta_j} \right),$$

where  $\alpha = (\alpha_{d+1}, \dots, \alpha_n)$ ,  $\alpha_{d+1}, \dots, \alpha_n \geq 0$ . Since  $|h_\alpha(\zeta_{d+1}, \dots, \zeta_n)| \leq h_\alpha(|\zeta_{d+1}|, \dots, |\zeta_n|)$ , it is enough to find an  $\alpha$  such that

$$\mathbb{E}' \cap \mathbb{R}_+^{n-d} \ni (t_{d+1}, \dots, t_n) \mapsto h_\alpha(t_{d+1}, \dots, t_n) r_d^{q_d}(t)$$

attains its maximum at  $(z_{d+1}, \dots, z_n)$ . Considering the partial (logarithmic) derivatives results in the following equations:

$$\begin{aligned} 0 &= \frac{1}{t_j} + \alpha_j - 2p_j q_d \frac{t_j^{2p_j-1}}{r_d(t)}, \quad j = d+1, \dots, k, \\ 0 &= \alpha_j - 2p_j q_d \frac{t_j^{2p_j-1}}{r_d(t)}, \quad j = k+1, \dots, n. \end{aligned}$$

We continue as in the proof of Proposition 2.3. □

*Proof of Proposition 2.5.* Consider the following two lemmas.

**Lemma 4.1.** *Assume that there exist  $0 < c < b < 1$  such that the function*

$$\varphi : [0, b] \ni t \longmapsto (1 - t^p)^{-q}$$

*is strictly concave and*

$$\varphi(0) + \frac{b}{c}(\varphi(c) - \varphi(0)) > \varphi(b) + 2.$$

*Let  $f \in \mathcal{O}(\mathbb{D})$ ,  $f \not\equiv 0$  such that  $|f(\zeta)|/\varphi(|\zeta|)$  attains its maximum at  $w_0$ . Then  $w_0 = 0$  or  $|w_0| \geq c$ .*

*Proof.* Assume that  $|w_0| \in (0, c)$ . We may assume that  $|f(w_0)| = \varphi(|w_0|)$ . Consider the function

$$\psi : [0, b] \ni t \longmapsto |f(0)| + \frac{t}{|w_0|}|f(w_0) - f(0)|.$$

From  $\psi(0) \leq \varphi(0)$ ,  $\psi(|w_0|) \geq \varphi(|w_0|)$ , and the convexity condition we get:

$$\begin{aligned} \psi(b) &= |f(0)| + \frac{b}{|w_0|}|f(w_0) - f(0)| \\ &\geq \varphi(0) + \frac{b}{|w_0|}|\varphi(|w_0|) - \varphi(0)| \\ &\geq \varphi(0) + \frac{b}{c}|\varphi(c) - \varphi(0)| > \varphi(b) + 2. \end{aligned}$$

The Schwarz lemma and the maximum principle imply that there exists a  $w \in \mathbb{D}$  with  $|w| = b$  and

$$\frac{|f(w) - f(0)|}{|w|} \geq \frac{|f(w_0) - f(0)|}{|w_0|}.$$

This means that

$$\begin{aligned} |f(w)| &\geq |f(w) - f(0)| - |f(0)| = |f(0)| + |f(w) - f(0)| - 2|f(0)| \\ &\geq \psi(b) - 2|f(0)| > \varphi(b) + 2 - 2|f(0)| \geq \varphi(b) = \varphi(|w|). \end{aligned}$$

This contradicts the maximality of  $|f(\zeta)|/\varphi(|\zeta|)$  at  $w_0$ .  $\square$

**Lemma 4.2.** *Assume that  $p \in (0, 1)$ ,  $q > 0$ . Then there exist  $b \in (0, 1)$ ,  $c \in (0, b)$  as required in Lemma 4.1.*

*Proof.* We have

$$\varphi'(t) = \frac{pq}{t^{1-p}(1-t^p)^{q+1}}$$

which is decreasing for small  $t$ . To prove the existence of  $c$  observe that

$$\lim_{d \rightarrow 0^+} \frac{\varphi(d) - \varphi(0)}{d} = +\infty.$$

$\square$

Observe that  $d = k$  for  $|z_\ell|$  small enough,  $\ell = 1, \dots, k$ . Let  $h \in \mathcal{O}(\mathbb{E}')$  and consider  $f(\zeta) := h(\zeta, 0, \dots, 0)$ ,  $\zeta \in \mathbb{D}$ . Put  $p := 2p_{k+1}$ ,  $q = q_d$ . Let  $b, c$  be as in Lemma 4.2. It follows from Lemma 4.1 that  $0 \neq |w_0| < c$  cannot be a maximum of  $\mathbb{D} \ni \zeta \mapsto |f(\zeta)|/\varphi(|\zeta|)$ . In particular  $z$  cannot be a maximum of  $\mathbb{E}' \ni \zeta \mapsto |h(\zeta)|r_d^{q_d}(\zeta)$  for  $0 \neq |z_{k+1}| < c$ .  $\square$

*Proof of Proposition 2.6.* Consider the following lemma.

**Lemma 4.3.** *Let  $a, c > 0$ ,  $t_0 \in (0, 1)$  be such that  $c \geq 1$  or  $4a + 2c > 1 + c^2$ ,  $t_0^c > \tau := a/(a + c)$ . Then there exist  $b > 0$  and  $r \geq 1$  such that*

$$h : [0, 1] \ni t \mapsto \frac{t^a}{(r - t)^b}(1 - t^c) \in [0, 1]$$

*admits its maximum at  $t_0$ .*

*Proof.* Comparing  $\varphi := th'(t)/h(t)$  to zero we get the equation:

$$\varphi(r, t) = a + \frac{bt}{r - t} - \frac{ct^c}{1 - t^c} = 0.$$

This gives us a formula for  $r$ :

$$r(t) = \frac{t(b - bt^c - a + at^c + t^c c)}{-a + at^c + t^c c}.$$

Observe that:

$$\lim_{t \rightarrow \tau^{1/c}} r(t) = +\infty,$$

$$\lim_{t \rightarrow 1} r(t) = 1.$$

In order to prove that  $r(t) > 1$  and that  $\varphi(r, \cdot)$  has only one zero it suffices to show that  $r'(t) < 0$ . We have:

$$r'(t) = \frac{(-(a + c)t^{2c} + (2a + c - c^2)t^c - a)b + (at^c + t^c c - a)^2}{(at^c + t^c c - a)^2}.$$

It remains to show that the coefficient  $\alpha(t^c)$  next to  $b$  is negative. We have:

$$\alpha(\tau) = \frac{-ac^2}{a + c} < 0,$$

$$\alpha(1) = -c^2 < 0,$$

$$\alpha'(u) = -2(a + c)u + 2a + c - c^2.$$

Let  $u_0$  be the zero of  $\alpha'(u)$ . For  $c \geq 1$  we have  $u_0 \leq \tau$  and we are done. Otherwise  $u_0 \in (\tau, 1)$  and  $4(a + c)\alpha(u_0) = c^2(1 + c^2 - 4a - 2c) < 0$ .  $\square$

We may assume that  $z_0 > 0$ . Put  $a = 2p_1$ ,  $c = 2p_2$ ,  $t_0 = z_0$ . Let  $r$  be as in Lemma 4.3. Putting

$$h(\zeta) = \frac{\zeta}{(r - \zeta)^{b/a}}.$$

completes the proof.  $\square$



**Remark 4.4.** The calculations in the proof of Proposition 2.6 can be performed using alternative function families e.g.

$$\begin{aligned} h_j(\zeta) &:= (\zeta + r)^k, \quad r \geq 0, \quad k = 0, 1, 2, \dots, \\ h_j(\zeta) &:= (\zeta + 1)^\alpha, \quad \alpha > 0, \\ h_j(\zeta) &:= (r - \zeta)^\alpha \zeta^k, \quad \alpha < 0, r \geq 1, k = 0, 1, 2, \dots \\ h_j(\zeta) &:= \left( \frac{\zeta + \delta}{1 + \delta\zeta} \right) \zeta^k, \quad \delta \in [0, 1], \quad k = 0, 1, 2, \dots \end{aligned}$$

However, the author was unable to solve the general case using any of them.

**Acknowledgement.** The author is indebted to professor Peter Pflug for his valuable comments on the paper.

#### REFERENCES

- [Jar-Jar-Pfl 2003] M. Jarnicki, W. Jarnicki, P. Pflug, *On extremal holomorphically contractible families*, Ann. Polon. Math. (2003), to appear.
- [Jar-Pfl 1993] M. Jarnicki, P. Pflug, *Invariant Distances and Metrics in Complex Analysis*, de Gruyter Exp. Math. 9, de Gruyter, 1993.
- [Lár-Sig 1998] F. Lárusson, R. Sigurdsson, *Plurisubharmonic functions and analytic discs on manifolds*, J. Reine Angew. Math. 501 (1998), 1–39.
- [Ngu 2003] Nguyen Quang Dieu, *Continuity of pluricomplex Green functions with poles along a hypersurface*, preprint.

JAGIELLONIAN UNIVERSITY, INSTITUTE OF MATHEMATICS  
REYMONTA 4, 30-059 KRAKÓW, POLAND

*E-mail address:* `wmj@im.uj.edu.pl`

*Current address:* Universität Osnabrück, Fachbereich Mathematik/Informatik  
Albrechtstraße 28, 49069 Osnabrück, Germany